

Literatur

1. Lippi G, von Meyer A, Cadamuro J, Simundic A-M. Blood sample quality. *Diagnosis*. 2018;6(1):25–31. doi:10.1515/dx-2018-0018.
2. O'Hara M, Wheatley EG, Kazmierczak SC. The impact of undetected in vitro hemolysis or sample contamination on patient care and outcomes in point-of-care testing: a retrospective study. *J Appl Lab Med*. 2020;5(2):332–341. doi:10.1093/jalm/jfz020.
3. Lippi G, Salvagno GL, Favaloro EJ, Guidi GC. Survey on the prevalence of hemolytic specimens in an academic hospital according to collection facility: opportunities for quality improvement. *Clin Chem Lab Med*. 2009;47(5):616–618. doi:10.1515/CCLM.2009.132.
4. Phelan MP, Hustey FM, Good DM, Reineks EZ. Seeing red: blood sample hemolysis is associated with prolonged emergency department throughput. *J Appl Lab Med*. 2020;5(4):732–737. doi:10.1093/jalm/jfaa073.
5. Wilson M, Adelman S, Maitre JB, et al. Accuracy of hemolyzed potassium levels in the emergency department. *West J Emerg Med*. 2020;21(6):272–275. doi:10.5811/westjem.2020.8.46812.
6. Milutinović D, Andrijević I, Ličina M, Andrijević L. Confidence level in venipuncture and knowledge on causes of in vitro hemolysis among healthcare professionals. *Biochem Med*. 2015;25(3):401–409. doi:10.11613/BM.2015.040.
7. Phelan MP, Ramos C, Walker LE, et al. The hidden cost of hemolyzed blood samples in the emergency department. *J Appl Lab Med*. 2021;6(6):1607–1610. doi:10.1093/jalm/jfab035.
8. Nichols JH, Apple FS. Prevalence of hemolyzed results in acute care settings. *J Appl Lab Med*. 2023;8:431–434. doi:10.1093/jalm/jfac141.
9. Tóth J, Oláh AV, Petercsák T, et al. Detection of haemolysis, a frequent preanalytical problem in the serum of newborns and adults. *EJIFCC*. 2020;31(1):6–14.
10. Khedr S, Blake V, Erdogan E. Neonatal unit hemolysis rates from an academic medical center: A quality improvement project. *Arch Path Lab*. 2016;140(6):502–503. doi:10.5858/arpa.2015-0252-le.
11. Lippi G, Fontana R, Avanzini P, Sandei F, Ippolito L. Influence of spurious hemolysis on blood gas analysis. *Clin Chem Lab Med*. 2013;51(8):1651–1654. doi:10.1515/cclm-2012-0802.
12. Hawkins R. Measurement of whole-blood potassium—is it clinically safe? *Clin Chem*. 2003;49(12):2105–2106. doi:10.1373/clinchem.2003.027227.
13. Werfen. GEM Premier 7000 with iQM3 Operators Manual. P/N 0000029366. Rev00. Nov 2023.
14. Dietzen DJ, Jackups R, Zaydman MA. Clinical implications of inaccurate potassium determination in hemolyzed pediatric blood specimens. *Clin Chim Acta*. 2024;15(557):117862. doi:10.1016/j.cca.2024.117862.
15. Wu AHB, Peacock WF. Potential medical impact of unrecognized in vitro hypokalemia due to hemolysis: a case series. *Clin Chem Lab Med*. 2024;aop:1–5. <https://doi.org/10.1515/cclm-2024-0351>.

16. Azman WNW, Omar J, Koon TS, Ismai TST. Hemolyzed specimens: major challenge for identifying and rejecting specimens in clinical laboratories. *Oman Med J*. 2019;34(2):94–98.
17. Möckel M, Luppä PB. Why hemolysis detection should be an integral part of any near-patient blood gas analysis. *J Lab Med*. 2021;45(4-5):193–195.
<https://doi.org/10.1515/labmed-2021-0076>.
18. Nigro M, Valli G, Marchionne ML, et al. Is there a risk of misinterpretation of potassium concentration from undetectable hemolysis using a POCT blood gas analyzer in the emergency department? *Medicina*. 2023;59(66):1–11.
<https://doi.org/10.3390/medicina59010066>.
19. Centers for Disease Control and Prevention. National hospital ambulatory medical care survey: 2021 emergency department summary tables. Accessed August 21, 2024. Available at
https://www.cdc.gov/nchs/data/nhamcs/web_tables/2021-nhamcs-ed-web-tables-508.pdf.